



Managed by Fermi Research Alliance, LLC for the U.S. Department of Energy Office of Science

Muon Physics Working Group Summary & Questions Posed

Andrew Norman (FNAL)

Angela Papa (PSI)

Haibo Li (IHEP, Beijing)

NuFact 2015 Rio de Janeiro, Brazil

Look Back to NuFact 13/14

Questions moving forward for 2014

- Three “Big” questions we want to address:

- Expt: What is the ultimate $\mu \rightarrow e\gamma$ and $\mu \rightarrow eee$ reach once $\mu N \rightarrow e N$ has set the limit.
 - What are the roles of the ratios of cLFV processes and other precision experiments at this point?
- Beams: What are the beam specifications for muon physics? (our requirements)
 - Are these compatible with the NuFact?
 - Are there other options?
- Theory: What else besides cLFV? EDMs?
 - What does theory tell us once we observe cLFV?
 - How do we relate our results to the models?

We understand the experiment design

- This generation
- And next

Many optimized designs that are starting into construction

- g-2, Mu2e, COMET, MEG2, Mu3e, DeeMe, etc...

Look Back to NuFact 13/14

Questions moving forward for 2014

- Three “Big” questions we want to address:
 - Expt: What is the ultimate $\mu \rightarrow e\gamma$ and $\mu \rightarrow eee$ reach once $\mu N \rightarrow eN$ has set the limit.
 - What are the roles of the ratios of cLFV processes and other precision experiments at this point?
 - Beams: What are the beam specifications for muon physics? (our requirements)
 - Are these compatible with the NuFact?
 - Are there other options?
 - Theory: What else besides cLFV? EDMs?
 - What does theory tell us once we observe cLFV?
 - How do we relate our results to the models?

We know the beams we need:

- need highly sculpted pulsed beams (Mu2e, COMET)
- need DC beams (MEG, Mu3e)
- need ultra cold beams (J-PARC g-2)

These MAY be compatible with a new facility

- Cost

May be options

- cooled μ 's + phase rot.

Questions Posed to NuFact 2015

- With the next generation of precision measurements experiments starting (muon g-2, mu to e conversion) are there additional measurement which can be made to ensure their success or to improve their background estimates and expand their sensitivities?
- Are there connections between cLFV searches and precision measurements which can be exploited to improve our progress in both regimes?
- How could a neutrino factory (and the supporting accelerator complex) be exploited to push our sensitivities for cLFV searches and precision measurements? Is there a secondary physics program that could be supported by such a complex which could address questions in both HEP and Nuclear Physics?
- What is the global picture that ties together “direct” and “indirect” for BSM physics? What are the results from the LHC and B-Factories telling us and is there a way to connect this sector with the next generation of precision measurements?

Questions Posed to NuFact 2015

- With the next generation of precision measurements experiments starting (muon g-2, mu to e conversion) are there additional measurement which can be made to ensure their success or to improve their background estimates and expand their sensitivities?
- Are there connections between cLFV searches and precision measurements which can be exploited to improve our progress in both regimes?
- How could a neutrino factory (and the supporting accelerator complex) be exploited to push our sensitivities for cLFV searches and precision measurements? Is there a secondary physics program that could be supported by such a complex which could address questions in both HEP and Nuclear Physics?
- What is the global picture that ties together “direct” and “indirect” for BSM physics? What are the results from the LHC and B-Factories telling us and is there a way to connect this sector with the next generation of precision measurements?

Global Perspective

- What is the global picture that ties together “direct” and “indirect” for BSM physics?
 - What are the results from the LHC and B-Factories telling us?
 - How do we connect this sector’s limits with the next generation of precision measurements?
 - Time scales?


Hints from the Energy Frontier

Introduction

arXiv:1207.7235

- LFV couplings to the Higgs possible ...
 - ... if SM only valid to finite scale Λ
 - ... in models with > 1 Higgs doublet (e.g., 2HDM)
- LFV Higgs couplings would allow processes like $\mu \rightarrow e$, $\tau \rightarrow \mu$ and $\tau \rightarrow e$ via a virtual Higgs boson

$$Y_{ij} = \frac{m_i}{v} \delta_{ij} + \frac{v^2}{\sqrt{2}\Lambda^2} \hat{\lambda}_{ij}$$



- $\mathcal{B}(H \rightarrow e\mu) < O(10^{-8})$ @ 95% CL from searches for $\mu \rightarrow e\gamma$
- $\mathcal{B}(H \rightarrow e\tau) < O(10\%)$ and $\mathcal{B}(H \rightarrow \mu\tau) < O(10\%)$ @ 95% CL from searches for $\tau \rightarrow e/\mu\gamma$ and μ/e g-2 measurements
- $\mathcal{B}(H \rightarrow e/\mu\tau) < 13\%$ @ 95% CL from theoretical reinterpretation of $H \rightarrow \tau\tau$ search results from ATLAS
→ direct search very promising


Hints from the Energy Frontier

Introduction

arXiv:1207.7235

- LFV couplings to the Higgs possible ...
 - ... if SM only valid to finite scale Λ
 - ... in models with > 1 Higgs doublet (e.g., 2HDM)
- LFV Higgs couplings would allow processes like $\mu \rightarrow e$, $\tau \rightarrow \mu$ and $\tau \rightarrow e$ via a virtual Higgs boson

$$Y_{ij} = \frac{m_i}{v} \delta_{ij} + \frac{v^2}{\sqrt{2}\Lambda^2} \hat{\lambda}_{ij}$$




- $\mathcal{B}(H \rightarrow e\mu) < O(10^{-8})$ @ 95% CL from searches for $\mu \rightarrow e\gamma$
- $\mathcal{B}(H \rightarrow e\tau) < O(10\%)$ and $\mathcal{B}(H \rightarrow \mu\tau) < O(10\%)$ @ 95% CL from searches for $\tau \rightarrow e/\mu\gamma$ and μ/e g-2 measurements
- $\mathcal{B}(H \rightarrow e/\mu\tau) < 13\%$ @ 95% CL from theoretical reinterpretation of $H \rightarrow \tau\tau$ search results from ATLAS
→ direct search very promising

Hints from the Energy Frontier

Results *Excess in $H \rightarrow \mu\tau$ channel*

arXiv:1502:07400



$$B(H \rightarrow \mu\tau)_{\text{best fit}} = (0.84^{+0.39}_{-0.37})\%$$

$$B(H \rightarrow \mu\tau) < 1.51\% \text{ @ } 95\% \text{ CL}$$

$$\sqrt{|Y_{\mu\tau}|^2 + |Y_{\tau\mu}|^2} < 3.6 \cdot 10^{-3}$$

Hints from the Energy Frontier

Events with μ and hadronically decaying τ .

Use τ kinematics and missing E_T to correct for undetected ν .



Two signal regions

SR1: $m_T(\mu, E_T^{\text{miss}}) > 40$ GeV and

$m_T(\tau_{\text{had}}, E_T^{\text{miss}}) < 40$ GeV


SR2: $m_T(\mu, E_T^{\text{miss}}) < 30$ GeV and

$m_T(\tau_{\text{had}}, E_T^{\text{miss}}) < 60$ GeV

Dominant backgrounds are $Z/\gamma^* \rightarrow \tau\tau$ (SR2)
and $W + \text{jets}$ (SR1)

$\text{BR} < 1.85\%$ (95% CL)

Theory: $\text{BR} < \sim 10\%$ from
 $\tau \rightarrow \mu\gamma$ and $(g-2)_{e,\mu}$



No Confirmation from Atlas in $H \rightarrow \mu\tau$ channel

Limits from the Energy Frontier

No evidence in
Z' and Quantum Black Holes


LFV in Z Decays

Results

- Count events in window around Z mass: (91 ± 3) GeV
- Background prediction of 83 ± 9
- Events found in data: 87
- Use CL_s method to determine limit:


$$B(Z \rightarrow e\mu)_{\text{expected}} < (6.7^{+2.8}_{-2.0}) \cdot 10^{-7}$$

$$B(Z \rightarrow e\mu)_{\text{observed}} < 7.3 \cdot 10^{-7}$$



LFV Decays of Heavy Resonances and QBHs

Results




No evidence in
Z Decays

Limits from the Energy Frontier

$Z \rightarrow e\mu$

PRD 90, 072010 (2014)
arXiv: 1408.5774

No evidence in
 $Z \rightarrow e\mu$




Fit to background + signal.

$BR(Z \rightarrow e\mu) < 7.5 \times 10^{-7}$ (95% CL)

LEP: $BR < 1.7 \times 10^{-6}$ (95% CL)

Limit inferred from
 $\mu \rightarrow eee$: $BR < 10^{-12}$



Limits from the Energy Frontier

$$Z \rightarrow e\mu$$

PRD 90, 072010 (2014)
arXiv: 1408.5774



No evidence in
 $Z \rightarrow e\mu$

Fit to background + signal.

$$Z' \text{ or } \tilde{\nu} \rightarrow e\mu, e\tau, \text{ or } \mu\tau$$


PRL 115, 031801 (2015), arXiv: 1503.04430

No evidence in
 $Z' \rightarrow \text{leptons}$






High Pt, back-to-back,
opposite sign, different
flavor.

Assume neutrino in same
direction as τ .



Craig Blocker (Brandeis University)



Craig Blocker (Brandeis University)

NuFact15

8

Limits from the Energy Frontier

$$Z \rightarrow e\mu$$

PRD 90, 072010 (2014)
arXiv: 1408.5774

No evidence in
 $Z \rightarrow e\mu$


$$Z' \text{ or } \tilde{\nu} \rightarrow e\mu, e\tau, \text{ or } \mu\tau$$

PRL 115, 031801 (2015), arXiv: 1503.04430

No evidence in
 $Z' \rightarrow \text{leptons}$

Multileptons in RPV SUSY

No evidence in
Multilepton channels



Craig Blocker (Brandeis)

Craig Blocker

Craig Blocker (Brandeis University)

NuFact15

15

Limits from the Energy Frontier

$$Z \rightarrow e\mu$$

PRD 90, 072010 (2014)
arXiv: 1408.5774

No evidence in
 $Z \rightarrow e\mu$

$$Z' \text{ or } \tilde{\nu} \rightarrow e\mu, e\tau, \text{ or } \mu\tau$$

PRL 115, 031801 (2015), arXiv: 1503.04430

No evidence in
 $Z' \rightarrow \text{leptons}$

Multileptons in RPV SUSY

No evidence in
Multilepton channels

Black Hole \rightarrow lepton + jet

PRL 112, 091804 (2014)


No evidence in
QBH searches

....

Quantum black holes predicted in low-scale gravity theories.

Expected to conserve angular momentum, charge, color but not other SM quantities.

Search for $BH \rightarrow \ell + \text{jet}$.



Limits from the B-Factories




Tau Lepton Flavor Violation



Identical in flavor structure to μ LFV channels

$$Br(\tau \rightarrow \mu\gamma)_{SM} \propto \left(\frac{\delta m_\nu^2}{m_W^2} \right)^2 < 10^{-40}$$

(X.Pham, EPJC8 513(1999))



Enhanced branching fractions due to τ mass

Ratio of Tau LFV decay BF: discrimination of NP models
JHEP 0705, 013(2007), PLB54 252 (2002)

Compare 10^{-7} for SUSY prediction to $10^{-14} - 10^{-11}$ prediction for $\mu \rightarrow e\gamma$


| | SUSY+GUT (SUSY+Seesaw) | Higgs mediated | Little Higgs | non-universal Z' boson |
|--|---------------------------|-------------------------|--------------|---------------------------|
| $\left(\frac{\tau \rightarrow \mu\mu\mu}{\tau \rightarrow \mu\gamma} \right)$ | $\sim 2 \times 10^{-3}$ | 0.06~0.1 | 0.4~2.3 | ~ 16 |
| $\left(\frac{\tau \rightarrow \mu ee}{\tau \rightarrow \mu\gamma} \right)$ | $\sim 1 \times 10^{-2}$ | $\sim 1 \times 10^{-2}$ | 0.3~1.6 | ~ 16 |
| $Br(\tau \rightarrow \mu\gamma)$ | $< 10^{-7}$ | $< 10^{-10}$ | $< 10^{-10}$ | $< 10^{-9}$ |

@Max

C. Cecchi

4

Limits from the B-Factories




$\tau \rightarrow 3 \text{ leptons}$



No evidence in
 $\tau \rightarrow 3\ell$

BELLE data set: 782 fb⁻¹
No events have been
found in the signal region

Very good lepton ID →
almost no bckgnd
Expected bckgnd events :
0.01 - 0.21




LHCb: 1 fb⁻¹@7TeV + 2fb⁻¹@8TeV
Br < 4.6 × 10⁻⁸ @ 90% C.L.
JHEP 02 (2015) 121

Br < (1.5-2.7) × 10⁻⁸ @ 90% C.L.
(Phys. Lett. B687, 139 (2010))

| Mode | ε (%) | $N_{\text{BG}}^{\text{EXP}}$ | σ_{sys} (%) | UL (×10 ⁻⁸) |
|---------------------|-------------------|------------------------------|---------------------------|-------------------------|
| $e^-e^+e^-$ | 6.0 | 0.21 ± 0.15 | 9.8 | 2.7 |
| $\mu^- \mu^+ \mu^-$ | 7.6 | 0.13 ± 0.06 | 7.4 | 2.1 |
| $e^- \mu^+ \mu^-$ | 6.1 | 0.10 ± 0.04 | 9.5 | 2.7 |
| $\mu^- e^+ e^-$ | 9.3 | 0.04 ± 0.04 | 7.8 | 1.8 |
| $\mu^- e^+ \mu^-$ | 10.1 | 0.02 ± 0.02 | 7.6 | 1.7 |
| $e^- \mu^+ e^-$ | 11.5 | 0.01 ± 0.01 | 7.7 | 1.5 |

Limits from the B-Factories

Or any of 48 other decay channels..... (but look at the sensitivities!)



48 decay modes investigated - 100 x more sensitivity w.r.t. CLEO results


LHCb results on 3 leptons comparable to B-factories

Limits from the Energy Frontier

- No clear evidence for LFV across MANY channels
- Current μ LFV experiments already set limits on many channels beyond the LHC reach
- But...many channels are not constrained.

Future at the Energy Frontier



- 14 TeV



Drives the cross sections and all the sensitivities to BSM physics

Future of the B-Factories

- ✓ SuperKEKB construction is finished
- ✓ startup for Phase 1 are in progress.



The no-background regime improves as $1 / \int L dt$
 If there are background events, the improvement is $1 / \sqrt{\int L dt}$

$\tau \rightarrow \mu \gamma$ (no bckgnd free) expected limite $O(10^{-9})$
 $\tau \rightarrow \mu \mu \mu$ (bckgnd free) expected limit $O(10^{-10})$

Questions Posed to NuFact 2015

- With the next generation of precision measurements experiments starting (muon g-2, mu to e conversion) are there additional measurement which can be made to ensure their success or to improve their background estimates and expand their sensitivities?
- Are there connections between cLFV searches and precision measurements which can be exploited to improve our progress in both regimes?
- How could a neutrino factory (and the supporting accelerator complex) be exploited to push our sensitivities for cLFV searches and precision measurements? Is there a secondary physics program that could be supported by such a complex which could address questions in both HEP and Nuclear Physics?
- What is the global picture that ties together “direct” and “indirect” for BSM physics? What are the results from the LHC and B-Factories telling us and is there a way to connect this sector with the next generation of precision measurements?


$\mu \rightarrow e\gamma$ (MEG and MEG2)

Results

Phy. Rev. Lett. 110, 201801 (2013)

Data taking finished at 31.08.2013

Statistics is doubled compare to published



| year | Nstop μ , $\times 10^{13}$ | Sensitivity, $\times 10^{-13}$ | Br, Upper limit (CL 90%), $\times 10^{-13}$ |
|---------------------|--------------------------------|--------------------------------|---|
| 2009+2010 | 17.5 | 13 | 13 |
| 2011 | 18.5 | 11 | 6,7 |
| 2009+2010+2011 | 36.0 | 7.7 | 5.7 (20 times better |
| All data (expected) | ~80 | ~5 | than MEGA) |


Final result of analysis is expected by the end of 2015 with the improved analysis. The data are reprocessed now.

$\mu \rightarrow e \gamma$ (MEG and MEG2)

Results

Phy. Rev. Lett. 110, 201801 (2013)

Data taking finished at 31.08.2013
Statistics is doubled compare to published



| year | Nstop μ , $\times 10^{13}$ | Sensitivity, $\times 10^{-13}$ | Br, Upper limit (CL 90%), $\times 10^{-13}$ |
|---------------------|--------------------------------|--------------------------------|---|
| 2009+2010 | 17.5 | 13 | 13 |
| 2011 | 18.5 | 11 | 6,7 |
| 2009+2010+2011 | 36.0 | 7.7 | 5.7 |
| All data (expected) | ~80 | ~5 | tha |

Final result of analysis is expected by the end of
with the improved analysis. The data are repro

Trying to improve the analysis
techniques to get better
sensitivity scaling than just \sqrt{N}
as they double their data set

Improvement of the analysis

- Event reconstruction algorithm.
- Calibration procedures.
- Background rejection techniques.
 - recover positron tracks which cross the target twice (missing turn analysis)
 - Identify background γ -rays generated when a positron annihilates with an electron on some detector material (annihilation-in-flight (AIF) analysis)
 - refine the alignment procedure of the target and drift chamber system.

$\mu \rightarrow e \gamma$ (MEG and MEG2)

How the sensitivity can be pushed down?

- More sensitive to the **signal**...

high statistics

$$SES = \frac{1}{R \times T \times A_g \times \epsilon(e^+) \times \epsilon(\text{gamma}) \times \epsilon(\text{TRG}) \times \epsilon(\text{sel})}$$

beam rate
acquisition time
geometrical acceptance
detector efficiency
selection efficiency

- More effective on rejecting the **background**...

high resolutions

$$B_{\text{acc}} \sim R \times \Delta E_e \times (\Delta E_{\text{gamma}})^2 \times \Delta T_{\text{egamma}} \times (\Delta \Theta_{\text{egamma}})^2$$


momentum resolution
Energy resolution
Relative timing resolution
Relative angular resolution

11

MEG2 focuses on improving energy and angular resolutions along with detector efficiencies to improve signal/background



- Redesigned spectrometer
- Upgraded calorimeter

The MEGII experiment -3D view



$\mu \rightarrow e\gamma$ (MEG and MEG2)

The upgraded Liquid Xenon calorimeter





- Calorimeter \rightarrow SiPM
- Higher density/eff dection

Re-designed timing counters
lower timing res to 35 ps for
pileup rejection

Goal is a 10x improvement in
sensitivities to $BR \sim 5 \times 10^{-14}$

A new re-designed spectrometer: the
pixelized Timing Counter (in numbers)

Timing resolution:
35 ps at the MEGII rate
conditions




$\mu \rightarrow eee$ (Mu3e)

The Mu3e Experiment

- Mu3e is a dedicated experiment searching for $\mu^+ \rightarrow e^+ e^- e^+$
- aimed sensitivity $\mathcal{B}(\mu \rightarrow eee) < 10^{-16}$
- stopped muons per second: 10^9
- main background: $\mu \rightarrow eee \nu_e \nu_\mu$, with $\mathcal{B} = 3.4 \cdot 10^{-5}$ and accidentals

Signal:




- $\Sigma \vec{p}_i = 0$
- common vertex
- $p < 53 \text{ MeV}$

BG: Internal Conversion



- $\Sigma \vec{p}_i \neq 0$
- common vertex


BG: Accidental



- non common vertex
- not in coincidence


Aug. 11, 2015 R. Gredig, NuFact15 Rio

Page 4




Phase 1a Detector Design

Beamline Design



- muon beamline at PSI
- low energy DC beams
- $\pi E5$ beamline: $\sim 10^8$ 28 MeV surface muons
 - shared with MEG
 - works for Phase I \rightarrow approved
- HiMB: High intensity muon beam study ongoing needed for Phase II
- Mu3e: $\sim 10^9 \mu^+/\text{s}$
- PSI Goal: $\sim 10^{10} \mu^+/\text{s}$

Beamline Progress



- muon beamline at PSI
- low energy DC beams
- $\pi E5$ beamline: $\sim 10^8$ 28 MeV/c surface muons
 - shared with MEG
 - works for Phase I \rightarrow approved
- HiMB: High intensity muon beam study ongoing needed for Phase II
- Mu3e: $\sim 10^9 \mu^+/\text{s}$
- PSI Goal: $\sim 10^{10} \mu^+/\text{s}$
- staged approach and modular principle
 - Phase Ia: Sensitivity $\mathcal{B}(\mu \rightarrow eee) < 10^{-14}$ (2016)
 - Phase Ib: Sensitivity $\mathcal{B}(\mu \rightarrow eee) < 10^{-15}$ (2017)
 - Phase II : Sensitivity $\mathcal{B}(\mu \rightarrow eee) < 10^{-16}$ (2019)

$\mu \rightarrow e$ Coherent Conversion

- Three different experimental designs & approaches:
 - Mu2e
 - Comet
 - DeeMe

- Measuring:

$$\mathcal{R} = \frac{\Gamma(\mu^- N \rightarrow e^- N)}{\Gamma(\mu^- N(Z) \rightarrow \nu_\mu N(Z-1))}$$

$$\mu^\pm \rightarrow e^\pm \gamma$$

MEG at PSI


$$\mu^\pm \rightarrow e^\pm e^+ e^-$$

Mu3e at PSI

$$\mu^- A(Z, N) \rightarrow e^- A(Z, N)$$

COMET at JPARC
Mu2e at FNAL

Mono energetic e emission at
endpoint of DIO spectrum




$\mu \rightarrow e$ Coherent Conversion

- Three different experimental designs & approaches:
 - Mu2e
 - Comet
 - DeeMe

Probes to effective energy scales of 1000's of TeV for BSM physics

Highly complementary to $\mu \rightarrow e \gamma$

One of most effective probes for new physics




$\mu A \rightarrow e A$ Designs

- Mu2e
 - Advanced detector and beamline designs
 - Forms part of core muon campus at FNAL



- Sensitivities of 2.9×10^{-17}
- First data in 2021



$\mu A \rightarrow e A$ Designs

- Under construction now

Mu2e Groundbreaking
April 18, 2015

Building for our future




Exploring the Unknown



$\mu A \rightarrow e A$ Designs

- COMET
 - Phased approach to reach sensitivity of 3×10^{-17}
 - Early sensitivity at 3×10^{-15}
- Advanced detector designs
- Det. prototypes & some prod. components complete
- Phase-I data in 2018/19

COMET: Phase-I




The COMET Experiment, 10 Aug 2015

17

Ben Krikler: bek07@imperial.ac

COMET: Phase-II




The COMET Experiment, 10 Aug 2015

15

Ben Krikler: bek07@imperial.ac.uk







μ A \rightarrow e A Designs

- Under construction now



$\mu A \rightarrow e A$ Designs



- DeeMe
 - Different approach uses beamline/spectrometer to momentum select signal e^- 's
 - Designed to reach SES 2×10^{-14}
 - Gated MWPCs to suppress beam flash
 - Engineering run in 2016



Electric field Profile

JPS Meeting 28/March/2014 @ Tokai University Shonan Campus

8



ermilab

Muon g-2

- One of the strongest probes for BSM physics
- Two very different approaches
 - FNAL g-2 uses “magic momentum” technique
 - J-PARC g-2 uses ultra cold muons and no electric focusing

BNL E821/Fermi

Magic momentum ($p=3.1 \text{ GeV}/c$)

$$\vec{\omega} = -\frac{e}{m} \left[a_{\mu} \vec{B} + \frac{\eta}{2} \left(\vec{\beta} \times \vec{B} + \frac{\vec{E}}{c} \right) \right]$$




$P=3.1 \text{ GeV}/c$
 $T=1.45 \text{ T}$

J-PARC

no electric focusing

$$\vec{\omega} = -\frac{e}{m} \left[a_{\mu} \vec{B} + \frac{\eta}{2} \left(\vec{\beta} \times \vec{B} \right) \right]$$




$P=0.3 \text{ GeV}/c$
 $T=3.0 \text{ T}$

FNAL Muon g-2

- Upgrade of BNL 821 to take advantage of FNAL beam complex + detector upgrades.
 - Uses same magnet (moved to Fermilab)
 - 4x improvement in statistical precision (0.54 ppm \rightarrow 140 ppb)

Improvement over BNL 821



- New quadrupoles
- New kicker modules
- New absolute field calibration
- New trolley field calibration system
- New detectors to measure beam profiles
- New analysis algorithms
- More complete simulation framework
- New data acquisition hardware
- New data acquisition software
- New laser gain stabilization system






FNAL Muon g-2

- Tremendous progress in ring re-commissioning & detector development
- First results expected 2018



New fixed and mobile field mapping probes


After 14 years, the ring has been cooled to superconducting temperatures and partially energized; some inevitable teething problems have been fixed, and the cooling should begin again on Monday



Three multiplane straw tracker systems will reconstruct the time-dependent muon decay position within the ring



J-PARC Muon g-2

- Ultra cold muon beam technique
- Precision ~ 100 ppb





J-PARC Muon g-2

- Experiment enabled by cold muon accelerator designs.
- Permits small magnet with no efield



Storage magnet



- 4 super-conducting coils supply injection field (Br), focusing field and main field.
 - Main field: 3T with local uniformity of 1ppm by iron shimming.

Questions Posed to NuFact 2015

- With the next generation of precision measurements experiments starting (muon g-2, mu to e conversion) are there additional measurement which can be made to ensure their success or to improve their background estimates and expand their sensitivities?
- Are there connections between cLFV searches and precision measurements which can be exploited to improve our progress in both regimes?
- How could a neutrino factory (and the supporting accelerator complex) be exploited to push our sensitivities for cLFV searches and precision measurements? Is there a secondary physics program that could be supported by such a complex which could address questions in both HEP and Nuclear Physics?
- What is the global picture that ties together “direct” and “indirect” for BSM physics? What are the results from the LHC and B-Factories telling us and is there a way to connect this sector with the next generation of precision measurements?

Other Measurements

- Mulan (muon lifetime)
- Mucap (muon capture on proton)
- Alcap (muon capture on Al)
- PiBeta/Pen (precision pion decays, rad. muon decay)

- nEDM
- NA61 (hadron yields)

- All of these measurements provide either sensitive probes for new physics, improve knowledge of systematics/backgrounds do both


MuLan

- Precision muon lifetime measurement
- Extracts Fermi's constant

$$\frac{1}{\tau_\mu} = \frac{G_F^2 m_\mu^5}{192\pi^3} \left(1 + \underbrace{\Delta q^{(0)}}_{\text{Phase space}} + \underbrace{\Delta q^{(1)}}_{\text{First order corrections}} + \underbrace{\Delta q^{(2)}}_{\text{Second order corrections}} \right)$$

$$G_F^{\text{MuLan}} = 1.166\,378\,7(6) \times 10^{-5} \text{ GeV}^{-2}$$

0.5 ppm



MuLan was systematics limited ... could we do better at a future facility?

| Uncertainty | R06 (ppm) | R07 (ppm) |
|---------------------------------|--------------|--------------|
| Kicker stability | 0.20 | 0.07 |
| μ SR distortions | 0.10 | 0.20 |
| Pulse pileup | | 0.20 |
| Gain variations | | 0.25 |
| Upstream stops | | 0.10 |
| Timing pick-off stability | | 0.12 |
| Master clock calibration | | 0.03 |
| Combined systematic uncertainty | 0.42 | 0.42 |
| Statistical uncertainty | 1.14 | 1.68 |


MuLan Collaboration, Phys. Rev. Lett. 99, 032001 (2007)

My Verdict: Probably ...

Mucap

- Muon capture on the proton
- Precision test of Chiral Perturbation Theory
- Measures g_p

Precise and unambiguous MuCap result confirms chiral perturbation theory prediction



$$g_p(\text{MuCap}) = 8.14 \pm 0.55$$




Hadronic Current

$$J^\alpha = \bar{u}_n \left(\underbrace{g_V \gamma^\alpha + \frac{i g_M}{2m_N} \sigma^{\alpha\nu} q_\nu + \frac{g_S}{m_\mu} q^\alpha}_{V^\alpha} - \underbrace{g_A \gamma^\alpha \gamma_5 + \frac{g_P}{m_\mu} q^\alpha \gamma_5 + \frac{i g_T}{2m_N} \sigma^{\alpha\nu} q_\nu \gamma_5}_{A^\alpha} \right) u_p$$

- CVC + G-Parity
 - $g_S, g_T \approx 0$
- CVC + Electron scattering
 - $g_V(q_\mu^2) = 0.976 \pm 0.001$
- Neutron beta decay
 - $g_A(q_\mu^2) = 1.2497 \pm 0.004$
 - Propagate $g_A(0) = 1.2723 \pm 0.0023 \rightarrow q^2$
 - K.A. Olive et al. (Particle Data Group), Chin. Phys. C, 38, 090001 (2014).
- This leaves g_p
 - Known with $\approx 50\%$ uncertainty


- Use a low-energy muon beam
- Stop in a specially prepared pure hydrogen target
- Image the stopping muon (TPC)
- Measure the disappearance rate
- Compare to the positive muon lifetime (MuLan)



Alcap

- Direct measurement of muon capture rate on AL
- Normalization for $\mu \rightarrow e$ conversion experiments
- Joint Mu2e/Comet exp

Run-1: Setup



AlCap, 12 Aug 2015

11

Ben Krikler: bek07@imperial.ac.uk

Run-1: Results (on-going)

So far:


- Proton Emission Spectrum from 3.5 to 10 MeV
- Proton emission per muon capture:
 - 0.020 (from 4 to 8 MeV)
 - 0.035 (integrated extrapolation)
 - Uncertainty about 9%

On-going:

- Deuteron, triton and alpha bands
- Final, combined proton emission rate

Impacts:

- COMET: Proton absorber removed
- Mu2e: Proton absorber re-optimised
- Changes to simulation code:
 - Vanilla Geant4 predicts about 30% of muon captures produce a proton
 - Fluka similar



AlCap, 12 Aug 2015

18

Ben Krikler: bek07@imperial.ac.uk

30 MeV/c,
3-6 KHz

Run-2 under analysis
Run-3 scheduled for Nov. 2015

PiBeta & PEN

- Measurements of: $\pi_{e3}, \pi_{e2}, \pi_{e2\gamma}$ and radiative muon decay
- Precision tests of SM, CKM elements, lepton universality, BSM searches

Reach of π_{e2} decay beyond the SM (New Physics)

$$\mathcal{L}_{NP} = \left[\pm \frac{\pi}{2\Lambda_V^2} \bar{u}\gamma_\alpha d \pm \frac{\pi}{2\Lambda_A^2} \bar{u}\gamma_\alpha \gamma_5 d \right] \bar{e}\gamma^\alpha (1 - \gamma_5)\nu + \left[\pm \frac{\pi}{2\Lambda_S^2} \bar{u}d \pm \frac{\pi}{2\Lambda_P^2} \bar{u}\gamma_5 d \right] \bar{e}(1 - \gamma_5)\nu, \quad (\Lambda_i \dots \text{scale of NP})$$

CKM unitarity and superallowed Fermi nuclear decays currently limit:

$$\Lambda_V \geq 20 \text{ TeV}, \quad \text{and} \quad \Lambda_S \geq 10 \text{ TeV}.$$



At $\Delta R_{e/\mu}^\pi / R_{e/\mu}^\pi = 10^{-3}$, π_{e2} decay is directly sensitive to:

$$\Lambda_P \leq 1000 \text{ TeV} \quad \text{and} \quad \Lambda_A \leq 20 \text{ TeV},$$

and indirectly, through loop effects to $\Lambda_S \leq 60 \text{ TeV}$.

The PIBETA/PEN apparatus

- $\pi E1$ beamline at PSI
- stopped π^+ beam
- active target counter
- 240-detector, spherical pure CsI calorimeter
- central tracking
- beam tracking
- digitized waveforms
- stable temp./humidity




RMD preliminary results, cont'd.

Preliminary result for RMD branching ratio (thesis E. Munyangabe):

$$B_{\text{exp}} = 4.365(9)_{\text{stat.}}(42)_{\text{syst.}} \times 10^{-3}, \quad \boxed{29 \times}$$

$$B_{\text{SM}} = 4.342(5)_{\text{stat-MC}} \times 10^{-3} \quad (\text{for } E_\gamma > 10 \text{ MeV}, \theta_{e\gamma} > 30^\circ)$$



Analysis of PS subset:


$13 \text{ MeV} < E_\gamma < 45 \text{ MeV}$, and
 $10 \text{ MeV} < E_{e^+} < 43 \text{ MeV}$, yields

$$\bar{\eta} = 0.006(17)_{\text{stat.}}(18)_{\text{syst.}}, \quad \text{or} \\ \bar{\eta} < 0.028 \quad (68\% \text{CL}).$$

$\sim 4 \times$ better than best previous experiment (Eichenberger et al, 84).

NB: preliminary results!

Neutron EDM



[2] Baker et al., PRL 97 (2006) 131801

Excellent (but beam limited) probe for BSM physics

Our apparatus is functioning well:

- Sensitivity is excellent
- Systematic effects are under control $< 5 \times 10^{-27} e \cdot \text{cm}$

We should reach $1.5 \times 10^{-26} e \cdot \text{cm}$ by mid 2016!

Next stage is to build a new setup (n2EDM) which should be able to reach $3 \times 10^{-27} e \cdot \text{cm}$

Questions Posed to NuFact 2015

- With the next generation of precision measurements experiments starting (muon g-2, mu to e conversion) are there additional measurement which can be made to ensure their success or to improve their background estimates and expand their sensitivities?
- Are there connections between cLFV searches and precision measurements which can be exploited to improve our progress in both regimes?
- How could a neutrino factory (and the supporting accelerator complex) be exploited to push our sensitivities for cLFV searches and precision measurements? Is there a secondary physics program that could be supported by such a complex which could address questions in both HEP and Nuclear Physics?
- What is the global picture that ties together “direct” and “indirect” for BSM physics? What are the results from the LHC and B-Factories telling us and is there a way to connect this sector with the next generation of precision measurements?

Looking Forward

- Muon physics program is transitioning in the questions being asked
 - No longer design questions
 - More complementarity/synergy questions with existing & future programs/facilities
- Starting new era where a convergence of new results is on horizon
 - LHC 14 TeV, Super-B Factories, g-2 and phase 1 mu2e's and nEDMs all may have new information in the next 5 year
 - May be able to start strongly constraining the BSM sector

New Questions

- Neutrino/Muon Physics:
 - “What overlaps exist in non-standard interactions? How would these manifest in both the near term muon/precision measurements sector & in the neutrino sector?”
- Beam/Machine Design:
 - “How can you improve experiments with out increasing the beam power? Cooled muon beams w/ phase rotations? New methods?”
- Program Planning:
 - “How do you support the physics needs for both DC and pulsed (high sculpted) beam structures in the planning (and cost) of new facilities?”

-
- Obrigado!
for a very successful NuFact!

